

Symmetry Analysis of Cylindrical Helmholtz Equation

Mehdi Nadjafikhah* Ali Mahdipour-Shirayeh†

Abstract

In this paper, we present the point symmetry group of three-dimensional homogeneous Helmholtz equation, when we consider the cylindrical coordinate system. In continuation, we present a complete set of functionally independent invariants of the equation along with the form of the general solution provided by these invariants. Finally, we find an optimal system of one-dimensional Lie subgroups of the full symmetry group.

A.M.S. 2000 Subject Classification: 34C14, 35J05, 70G65.

Keywords: Helmholtz equation, Lie point symmetries, optimal system of Lie subalgebras.

1 Introduction

The Helmholtz equation, named for Hermann von Helmholtz, is the elliptic partial differential equation (in homogenous case)

$$\nabla u + k^2 u = 0,$$

when u is a vector function and ∇ is the vector Laplacian.

The three-dimensional homogeneous *cylindrical Helmholtz equation* (in brief CHE), is a Helmholtz differential equation defined in the cylindrical (r, θ, z) coordinate system, i.e.

$$\text{CHE} : u_{rr} + \frac{1}{r}u_r + \frac{1}{r^2}u_{\theta\theta} + u_{zz} + k^2 u = 0, \quad (1)$$

where u is a function of variables r, θ, z and k is a constant.

The Helmholtz equation naturally appears from general conservation laws of physics and often arises in the study of physical problems involving partial differential equations (PDEs) in both space and time, more particularly in mechanics, acoustics, electromagnetics, etc. (see for details [3, 4, 5]).

*Department of Mathematics, Iran University of Science and Technology, Narmak-16, Tehran, I.R.Iran. e-mail: m_nadjafikhah@iust.ac.ir

†e-mail: mahdipour@iust.ac.ir

This equation can be interpreted as a wave equation for monochromatic waves and can be derived from the heat conduction equation, Schrödinger equation, telegraph and other wave type, or evolutionary equations [4]. When $k = 0$, the Helmholtz differential equation reduces to Laplace's equation and when $k^2 < 0$ (for imaginary), the equation becomes the space part of the diffusion equation.

In the mathematical sense, it appears as an eigenvalue problem for the Laplace operator ∇^2 (see [3, 4, 9]). As is well-known, there are some mathematical theory which can be applied to a solution of the Helmholtz equation in Cartesian, spherical, cylindrical coordinates, and etc., by applying the technique of separation of variables and boundary element method, its various solutions for many basic shapes, for example in a vibrating membrane, along with other different methods [1, 2, 4, 9]. In addition, since the Helmholtz equation is an elliptic equation, so it is usual to consider boundary value problem along with transmission conditions and integral representation of solutions (see [3, 4] and references therein).

At the present work, we are dealing with the symmetry analysis and a one-dimensional optimal system of the Helmholtz equation in cylindrical coordinate. Cylindrical coordinates are the natural choice for studying different shapes of the Helmholtz equation, that is a good motivation for work with this coordinate. Also we find a form of general solutions providing by the differential invariants of the equation.

2 Symmetry of CHE

In this section, we are deal with symmetry analysis of CHE, when we act the point symmetry group on the second order jet space $J^2 = J^2(\mathbb{R}^3, \mathbb{R})$ with local coordinates $(r, \theta, z, u, u^{(1)}, u^{(2)})$. In these charts $u^{(\alpha)}$ ($\alpha = 1, 2$) indicates the set of all α^{th} derivatives of the dependent variable u with respect to independent variable r, θ, z .

The base space of the jet bundle J^2 is the space $\mathcal{M} = X \times U$ of independent and dependent variables (x, u) for $x = (r, \theta, z)$, that is in fact the zero-order jet space $J^0(\mathbb{R}^3, \mathbb{R})$. CHE defines a subvariety of jet bundle J^2 of three-dimensional submanifold of $\mathcal{M}$, that is, graph of function $u = u(r, \theta, z)$.

The point symmetry group on $\mathcal{M}$ is defined by transformations in the general form of

$$\tilde{r} = \varphi(r, \theta, z, u), \quad \tilde{\theta} = \psi(r, \theta, z, u), \quad \tilde{z} = \chi(r, \theta, z, u), \quad \tilde{u} = \zeta(r, \theta, z, u),$$

for arbitrary smooth functions φ, ψ, χ and ζ .

Let

$$v = \sum_{i=1}^3 \xi^i(x, u) \frac{\partial}{\partial x_i} + \eta(x, u) \frac{\partial}{\partial u}, \quad (2)$$

be the general form of an infinitesimal generator of point symmetry transformations on $\mathcal{M}$, in which the coefficients are arbitrary smooth functions. Let the

second order prolongation of v be defined as

$$v^{(2)} = v + \sum_{\#J=j=1}^2 \hat{\eta}_J(x, u^{(j)}) \frac{\partial}{\partial u_J},$$

where we assumed that each index signifies to the derivation in respect to the index, and $J = (j_1, j_2)$ is a multi-index with length $0 \leq \#J \leq 2$. In addition, coefficients are given by the prolongation formula (see [10])

$$\hat{\eta}_J = D_J(Q) + \sum_{i=1}^3 \xi^i u_{J,i}.$$

In the latter relation, D_J is the total derivative with respect to the variables signified by J , and $Q = \eta - \sum_{i=1}^3 \xi^i u_i$ is the characteristic of v .

According to [10], the vector field v is an infinitesimal symmetry of Eq. (1), if and only if it satisfies the infinitesimal invariance condition

$$v^{(2)} \left(u_{rr} + \frac{1}{r} u_r + \frac{1}{r^2} u_{\theta\theta} + u_{zz} + k^2 u \right) = 0, \quad (3)$$

whenever Eq. (1) vanishes.

Eq. (3) leads to the infinitesimal determining equation, that must be solved for arbitrary functions ξ^i and η , when they are dependent to variables r, θ, z, u and not to the derivatives u in respect to its variables. Hence, the induced equation satisfies for ξ^i and η , if and only if the following determining equations are fulfilled

$$\begin{aligned} \xi_{uu}^1 &= 0, & \xi_{uu}^2 &= 0, & \xi_{uu}^3 &= 0, & \xi_u^2 &= 0, \\ \xi_{uu}^1 &= 0, & \xi_{uu}^3 &= 0, & \xi_u^1 &= 0, & \xi_u^1 &= 0, \\ \xi_u^2 &= 0, & \xi_u^1 &= 0, & \xi_u^3 &= 0, & \xi_u^3 &= 0, \\ r^2 \xi_z^2 + \xi_\theta^3 &= 0, & \xi_u^1 &= 0, & \xi_{uu}^3 &= 0, & \xi_z^1 + \xi_r^3 &= 0, \\ 2 \xi_u^1 + r \eta_{uu} - 2 r \xi_{ru}^1 &= 0, & r \xi_r^1 - r \xi_\theta^2 - \xi^1 &= 0, & \xi_r^1 - \xi_z^3 &= 0, \\ 2 \xi_{zu}^3 - \eta_{uu} &= 0, & r^2 \xi_r^2 + \xi_\theta^1 &= 0, & \xi_{ru}^3 + \xi_{zu}^1 &= 0, \\ \xi_{\theta u}^3 + r^2 \xi_{zu}^2 &= 0, & \eta_{uu} - 2 \xi_{\theta u}^2 &= 0, & r^2 \xi_{ru}^2 + \xi_{\theta u}^1 &= 0, \\ r^2 k^2 u \xi_u^2 + 2 \eta_{\theta u} - r \xi_r^2 - \xi_{\theta\theta}^2 - r^2 \xi_{rr}^2 - r^2 \xi_{zz}^2 &= 0, \\ 3 r^2 k^2 u \xi_u^1 + r \xi_r^1 + 2 r^2 \eta_{ru} - r^2 \xi_{zz}^1 - \xi^1 - \xi_{\theta\theta}^1 - r^2 \xi_{rr}^1 &= 0, \\ r^2 k^2 u \xi_u^3 + 2 r^2 \eta_{zu} - r \xi_r^3 - r^2 \xi_{zz}^3 - \xi_{\theta\theta}^3 - r^2 \xi_{rr}^3 &= 0, \\ r^2 \eta_{z,z} + 2 r^2 k^2 u (\xi_r^1 - \eta_u) + r^2 \eta_{rr} + r \eta_r + \eta_{\theta\theta} + r^2 \eta k^2 &= 0. \end{aligned} \quad (4)$$

The solution space $\mathfrak{g}$ of the system of determining equations (4) with respect to the unknown coefficients of v , is the Lie algebra of infinitesimal symmetry of

Eq. (1). By solving system (4) (that is applicable with MAPLE) we obtain the below solutions

$$\begin{aligned}
\xi^1 &= (c_1 z + c_3) \sin \theta + (c_2 z + c_4) \cos \theta, \\
\xi^2 &= -\frac{1}{r}(c_2 z + c_4) \sin \theta + \frac{1}{r}(c_1 z + c_3) \cos \theta + c_5, \\
\xi^3 &= -r(c_1 \sin \theta + c_2 \cos \theta) + c_6, \\
\eta &= \exp(-\sqrt{c_3}z - \sqrt{c_2}\theta)(c_7 \exp(2\sqrt{c_2}\theta) + c_8)(c_9 \exp(2\sqrt{c_3}z) + c_{10}) \\
&\quad \times \left(c_{11} \text{BY}(\sqrt{-c_2}, \sqrt{c_3 + k^2 r}) + c_{12} \text{BJ}(\sqrt{-c_2}, \sqrt{c_3 + k^2 r}) \right) + c_{13} u,
\end{aligned} \tag{5}$$

where, BJ and BY are the Bessel functions of the first and second kinds, resp. which satisfy Bessel's equation $x^2 y'' + x y' + (x^2 - v^2)y = 0$, and c_i (for $1 \leq i \leq 13$) are arbitrary constants.

Noting to the concluded form of infinitesimal operators of Eq. (1), one may divide the resulting form of v to the set of following vector fields, as a basis of Lie algebra $\mathfrak{g}$ of the symmetry group G of CHE

$$\begin{aligned}
X_1 &= \frac{\partial}{\partial \theta}, & X_2 &= \frac{\partial}{\partial z}, & X_3 &= u \frac{\partial}{\partial u}, \\
X_4 &= \sin \theta \frac{\partial}{\partial r} + \frac{\cos \theta}{r} \frac{\partial}{\partial \theta}, & X_5 &= -\cos \theta \frac{\partial}{\partial r} + \frac{\sin \theta}{r} \frac{\partial}{\partial \theta}, \\
X_6 &= -z \cos \theta \frac{\partial}{\partial r} + \frac{z \sin \theta}{r} \frac{\partial}{\partial \theta} + r \cos \theta \frac{\partial}{\partial z}, \\
X_7 &= z \sin \theta \frac{\partial}{\partial r} + \frac{z \cos \theta}{r} \frac{\partial}{\partial \theta} - r \sin \theta \frac{\partial}{\partial z}.
\end{aligned} \tag{6}$$

The commutator of every X_i and X_j i.e. $[X_i, X_j] := X_i X_j - X_j X_i$ ($1 \leq i, j \leq 7$), is straightforwardly a linear combination of vector fields (6). The complete set of commutators is given in Table 1.

	X_1	X_2	X_3	X_4	X_5	X_6	X_7
X_1	0	0	0	$-X_5$	X_4	X_7	$-X_6$
X_2	0	0	0	0	0	X_5	X_4
X_3	0	0	0	0	0	0	0
X_4	X_5	0	0	0	0	0	$-X_2$
X_5	$-X_4$	0	0	0	0	$-X_2$	0
X_6	$-X_7$	$-X_5$	0	0	X_2	0	X_1
X_7	X_6	$-X_4$	0	X_2	0	$-X_1$	0

Table 1: Commutator table for $\mathfrak{g}$.

Theorem 1. *The complete set of infinitesimal generators of point symmetry group of Eq. (1), is introduced in (2) and (5) and is a seven-dimensional Lie algebra with commutators given in Table 1.*

Since the derived algebra $\mathfrak{g}^{(1)} = [\mathfrak{g}, \mathfrak{g}] = \mathfrak{g} / \langle X_3 \rangle$ and for each $k > 1$, $\mathfrak{g}^{(k+1)} = [\mathfrak{g}^{(k)}, \mathfrak{g}^{(k)}] = \mathfrak{g}^{(1)}$, so the descending sequence of the derived subalgebras of $\mathfrak{g}$,

$$\mathfrak{g} \supset \mathfrak{g}^{(1)} = \mathfrak{g}^{(2)} = \dots$$

does not terminate with a null ideal and hence $\mathfrak{g}$ is not solvable [8, 12]. Also, $\mathfrak{g}$ is not semisimple, since by applying Cartan's criterion of semisimplicity [12], its Killing form defined for $v = \sum_i v_i X_i$ and $w = \sum_j w_j X_j$ in $\mathfrak{g}$ as

$$K(v, w) = \text{tr}(\text{ad}(v) \circ \text{ad}(w)) = -4(v_1 w_1 + v_6 w_6 + v_7 w_7)$$

is degenerate.

The radical of $\mathfrak{g}$ is equal to the Lie subalgebra $\mathfrak{r} := \langle X_2, X_3, X_4, X_5 \rangle$ and hence Levi decomposition of $\mathfrak{g}$ is in the form of $\mathfrak{g} = \mathfrak{r} \oplus \mathfrak{s}$, for semisimple subalgebra $\mathfrak{s} := \mathfrak{g} / \mathfrak{r} = \langle X_1, X_6, X_7 \rangle$. According to the Table 1, $\mathfrak{r}$ is an abelian group of dimension 4, thus is isomorphic to $\mathbb{R}^4$, and the (real) Lie subalgebra $\mathfrak{s}$ is isomorphic to the Lie algebra $\mathfrak{so}(3)$ of the special orthogonal group. In addition, $\mathfrak{s}$ is not an ideal of $\mathfrak{g}$ and therefore we find the structure of the symmetry algebra of CHE as following semi-direct product

$$\mathfrak{g} = \mathbb{R}^4 \ltimes \mathfrak{so}(3). \quad (7)$$

For finding the symmetry group G of the Lie algebra $\mathfrak{g}$ generated by infinitesimal operators (6), we determine the one-parameter group (flow) of these vector fields:

Theorem 2. *If we denote the group transformations of each infinitesimal generator X_i ($i = 1, \dots, 7$) with $g_i(s)$ for different values of parameter s , then we have*

$$\begin{aligned} g_1(s) &: (r, \theta, z, u) \mapsto (r, \theta + s, z, u), \\ g_2(s) &: (r, \theta, z, u) \mapsto (r, \theta, z + s, u), \\ g_3(s) &: (r, \theta, z, u) \mapsto (r, \theta, z, u + s), \\ g_4(s) &: (r, \theta, z, u) \mapsto \left(r \left(\cos^2 \theta + \frac{1}{4r^2} (2s - r \sin 2\theta)^2 \right)^{1/2}, \right. \\ &\quad \left. \arctan \left(\frac{s}{r} (1 + \tan^2 \theta)^{1/2} - \tan \theta \right), z, u \right), \\ g_5(s) &: (r, \theta, z, u) \mapsto \left(\left(s - \frac{r}{(1 + \tan^2 \theta)^{1/2}} \right) \left(1 + \left(\frac{r \tan \theta}{s(1 + \tan^2 \theta)^{1/2} - r} \right)^2 \right)^{1/2}, \right. \\ &\quad \left. \arctan \left(\frac{r \tan \theta}{s(1 + \tan^2 \theta)^{1/2} + r} \right), z, u \right), \end{aligned}$$

$$\begin{aligned}
g_6(s) &: (r, \theta, z, u) \longmapsto \left(\left(c_1 \cos^2(s + c_2) + d_1^2 \sin^2(s + c_2) \right)^{1/2}, \right. \\
&\quad \theta - \frac{\left((c_1 - d_1^2) \cos^2(s + c_2) \right)^{1/2} \tanh^{-1}\left(\frac{d_2^2 - c_1}{d_1} \cos(s + c_2) \right)}{(d_1^2 - c_1)^{1/2} \cos(s + c_2)} \\
&\quad + \frac{\left((c_1 - d_1^2) \cos^2(c_2) \right)^{1/2} \tanh^{-1}\left(\frac{(d_2^2 - c_1)}{d_1} \cos(c_2) \right)}{(d_1^2 - c_1)^{1/2} \cos(c_2)}, \\
&\quad \left. \frac{\sqrt{2}}{2} \left((c_1 - d_1^2) \sin(2s + 2c_2) \right)^{1/2}, u \right), \\
g_7(s) &: (r, \theta, z, u) \longmapsto \left(\left(c_1 \cos^2(s + c_3) + d_2^2 \sin^2(s + c_3) \right)^{1/2}, \right. \\
&\quad \theta + \frac{\left((c_1 - d_2^2) \cos^2(s + c_3) \right)^{1/2} \tanh^{-1}\left(\frac{d_2^2 - c_1}{d_1} \cos(s + c_3) \right)}{(d_2^2 - c_1)^{1/2} \cos(s + c_3)} \\
&\quad - \frac{\left((c_1 - d_2^2) \cos^2(c_3) \right)^{1/2} \tanh^{-1}\left(\frac{(d_2^2 - c_1)}{d_1} \cos(c_3) \right)}{(d_2^2 - c_1)^{1/2} \cos(c_3)}, \\
&\quad \left. \frac{\sqrt{2}}{2} \left((c_1 - d_2^2) \sin(2s + 2c_3) \right)^{1/2}, u \right),
\end{aligned}$$

where we assume that $c_1 = r^2 + z^2$, $c_2 = \arctan(z(r^2 \cos^2 t)^{-1/2})$, $c_3 = -s - \arctan(z(r^2 \sin^2 t)^{-1/2})$, $d_1 = r \sin t$ and $d_2 = r \cos t$.

It is well-known that for each one-parameter subgroup of the full symmetry group of a system there will correspond a family of solutions that are called invariant solutions [10, 11, 12]. In section four, we introduce a general solution of CHE with respect to its functionally independent invariants.

Theorem 3. *If $u = f(r, \theta, z)$ be a solution of (1), then are the below functions ($i = 4, \dots, 7$)*

$$\begin{aligned}
g_1(s) \cdot f(r, \theta, z) &= f(r, \theta + s, z), \\
g_2(s) \cdot f(r, \theta, z) &= f(r, \theta, z + s), \\
g_3(s) \cdot f(r, \theta, z) &= f(r, \theta, z) - s, \\
g_i(s) \cdot f(r, \theta, z) &= f(g_i(s) \cdot (r, \theta, z)),
\end{aligned}$$

when $g_i(s) \cdot (r, \theta, z)$ means the first three components of $g_i(s) \cdot (r, \theta, z, u)$ and s is an arbitrary parameter.

3 Invariants and General Solutions

In this section, we study the invariant solutions of various combinations of vector fields (6) based on the following method:

the invariants $u = I(r, \theta, z, u)$ of one-parameter group with infinitesimal generators in the form of (6) satisfy the linear homogeneous partial differential equations of first order:

$$v[I] = 0.$$

The solutions of the latter are found by the method of characteristics (See [6] and [11] for details). So we can replace the latest Eq. by the following characteristic system of ordinary differential equations

$$\frac{dr}{\xi_1} = \frac{d\theta}{\xi_2} = \frac{dz}{\xi_3} = \frac{du}{\eta}. \quad (8)$$

By solving the Eqs. (8) of differential generators (5), we (locally) find the following general solutions

$$\begin{aligned} I_1(r, \theta, z, u) &= 2z + \frac{(c_1 \sin \theta + c_2 \cos \theta) r^2 + c_6 r}{(c_1 z + c_3) \sin \theta + (c_2 z + c_4) \cos \theta} = d_1, \\ I_2(r, \theta, z, u) &= z - \frac{1}{c_{13}} (c_1 \sin \theta + c_2 \cos \theta) \ln \left((c_{11} \text{BY}(\sqrt{-c_2}, \sqrt{c_3 + k^2}) \right. \\ &\quad \left. + c_{12} \text{BJ}(\sqrt{-c_2}, \sqrt{c_3 + k^2})) (c_7 c_9 \exp(\sqrt{c_2} \theta + \sqrt{c_3} z) \right. \\ &\quad \left. + c_7 c_{10} \exp(\sqrt{c_2} \theta - \sqrt{c_3} z) + c_8 c_9 \exp(-\sqrt{c_2} \theta + \sqrt{c_3} z) \right. \\ &\quad \left. + c_8 c_{10} \exp(-\sqrt{c_2} \theta - \sqrt{c_3} z)) + c_{13} u \right) = d_2, \\ I_3(r, \theta, z, u) &= -\frac{1}{B} \arctan(\alpha) (2 c_5 A^{-1} (c_2 c_3 - c_1 c_4) r^3 + 2 c_6 r) \\ &\quad + r^2 A^{-1} \left(\ln(C) - \ln \left(\frac{1}{\cos \theta + 1} \right) \right) (c_2^2 z + c_1 c_3 + c_2 c_4 + c_1^2 z) \\ &\quad + 2 r^2 A^{-1} \arctan \left(\frac{\cos \theta - 1}{\sin \theta} \right) (c_2 c_3 - c_1 c_4) = d_3. \end{aligned} \quad (9)$$

when we assumed that

$$\begin{aligned} A &= c_2^2 z^2 + 2 c_2 c_4 z + c_4^2 + c_1^2 z^2 + 2 c_1 c_3 z + c_3^2 \\ B &= (-2 c_1 c_3 z - c_3^2 - c_1^2 z^2 + c_5^2 r^2 - c_2^2 z^2 - 2 c_2 c_4 z - c_4^2)^{\frac{1}{2}} \\ C &= \frac{1}{\cos \theta + 1} (c_1 z \cos \theta + c_3 \cos \theta - c_4 \sin \theta - c_2 z \sin \theta + c_5 r) \\ \alpha &= \frac{1}{B \sin \theta} (c_1 z - c_1 z \cos \theta + c_3 - c_3 \cos \theta - c_5 r + c_5 r \cos \theta + c_2 z \sin \theta + c_4 \sin \theta). \end{aligned}$$

and d_i s ($i = 1, 2, 3$) are some constants.

The functions I_1, I_2, I_3 form a complete set of functionally independent invariants of one-parameter group generated by (8) (see [10]).

According to the theorem of section 4.3.3 of [6], the derived invariants (9) as independent first integrals of the characteristic system of the infinitesimal generators (6), provide the general solution

$$S(r, \theta, z, u) := \mu(I_1(r, \theta, z, u), I_2(r, \theta, z, u), I_3(r, \theta, z, u)),$$

with an arbitrary function μ .

4 Classification of subalgebras

It is well-known that the problem of classifying invariant solutions is equivalent to the problem of classifying subgroups of the full symmetry group under conjugation in which itself is equivalent to determining all conjugate subalgebras [10, 12]. The latter problem, tends to determine a list (that is called an *optimal system*) of conjugacy inequivalent subalgebras with the property that any other subalgebra is equivalent to a unique member of the list under some element of the adjoint representation i.e. $\bar{\mathfrak{h}} \text{Ad}(g) \mathfrak{h}$ for some g of a considered Lie group.

For finding one-dimensional subalgebras, this classification problem is essentially the same as the problem of classifying the orbits of the adjoint representation.

The adjoint action is given by the Lie series

$$\text{Ad}(\exp(sX_i))X_j = X_j - s[X_i, X_j] + \frac{s^2}{2}[X_i, [X_i, X_j]] - \dots,$$

where s is a parameter and $i, j = 1, \dots, 7$. Let $M_i(s) : \mathfrak{g} \rightarrow \mathfrak{g}$ defined by $X \mapsto \text{Ad}(\exp(sX_i))X$ be the adjoint representation on $\mathfrak{g}$ (for $i = 1, \dots, 7$ and small parameter s) and also $M_i(s)$ be its matrix of representations with respect to the basis $\{X_1, \dots, X_7\}$, by an abuse of convention. After straightforwardly computations, we find that

$$\begin{aligned} M_1(s) &= \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \cos s & \sin s & 0 & 0 \\ 0 & 0 & 0 & -\sin s & \cos s & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \cos s & -\sin s \\ 0 & 0 & 0 & 0 & 0 & \sin s & \cos s \end{bmatrix}, \\ M_2(s) &= \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -s & 1 & 0 \\ 0 & 0 & 0 & -s & 0 & 0 & 1 \end{bmatrix}, \\ M_3(s) &= \text{Id}_7, \end{aligned}$$

$$\begin{aligned}
M_4(s) &= \begin{bmatrix} 1 & 0 & 0 & 0 & -s & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & s & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \\
M_5(s) &= \begin{bmatrix} 1 & 0 & 0 & s & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & s & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \\
M_6(s) &= \begin{bmatrix} \cos s & 0 & 0 & 0 & 0 & 0 & \sin s_6 \\ 0 & \cos s & 0 & 0 & \sin s & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & -\sin s & 0 & 0 & \cos s & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ -\sin s & 0 & 0 & 0 & 0 & 0 & \cos s \end{bmatrix}, \\
M_7(s) &= \begin{bmatrix} \cos s & 0 & 0 & 0 & 0 & -\sin s & 0 \\ 0 & \cos s & 0 & \sin s & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & -\sin s & 0 & \cos s & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ \sin s & 0 & 0 & 0 & 0 & \cos s & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}.
\end{aligned}$$

In the matrices M_1, M_6 and M_7 , there are minor matrices which indicate 3-dimensional rotations, that are related to the section $\mathfrak{so}(3)$ of the Lie structure (7).

Theorem 4. *An optimal system of one-dimensional Lie subalgebras of (1) is provided by those generated by*

- | | |
|----------------------|-------------------------------------|
| 1) X_1 , | 10) $a X_4 + b X_6$, |
| 2) X_2 , | 11) $a X_5 + b X_7$, |
| 3) X_6 , | 12) $X_3 + a X_1 + b X_2$, |
| 4) X_7 , | 13) $X_3 + a X_4 + b X_6$, |
| 5) $X_3 + a X_1$, | 14) $a X_1 + b X_2 + c X_5$, |
| 6) $X_3 + a X_4$, | 15) $a X_2 + b X_5 + c X_7$, |
| 7) $X_3 + a X_6$, | 16) $a X_4 + b X_5 + c X_6$, |
| 8) $X_3 + a X_7$, | 17) $X_3 + a X_1 + b X_2 + c X_5$, |
| 9) $a X_1 + b X_2$, | |

where the coefficients a , b and c are arbitrary constants.

Proof: Let $X = \sum_{i=1}^7 a_i X_i$ be an element of $\mathfrak{g}$, then we see that

$$\begin{aligned} M_7(s_7) \circ \cdots \circ M_2(s_2) \circ M_1(s_1) : X &\longmapsto \left(a_1 \cos s_6 \cos s_7 + \cdots + a_7 \sin s_6 \right) X_1 \\ &+ \left(a_2 \cos s_6 \cos s_7 + a_4 \cos s_6 \sin s_7 + a_5 \sin s_6 \right) X_2 \\ &\vdots \\ &+ \left(a_1 (\sin s_1 \sin s_7 - \cos s_1 \sin s_6 \cos s_7) + \cdots + a_7 \cos s_1 \cos s_6 \right) X_7. \end{aligned}$$

For classifying one-dimensional Lie subalgebras of CHE, we plan the following cases that in each case, by acting a finite number of the adjoint representations M_i ($i = 1, \dots, 7$) on X , by proper selection of parameters s_i in each stage, we gradually try to make the coefficients of X vanish and to acquire the most simple form of X .

- At first, we suppose that $a_3 \neq 0$ and scaling if necessary, we can assume that $a_3 = 1$. This assumption suggests *Case 1*, $\dots$, *Case 10*.

Case 1. If $a_5 \neq 0$, then we act on X by $\text{Ad}(\exp(-\arctan(a_4/a_5) X_1))$ and hence we can make the coefficient of X_4 vanish. Then we tend to the new form

$$X' = a_1 X_1 + a_2 X_2 + X_3 + a'_5 X_5 + a'_6 X_6 + a'_7 X_7$$

for certain scalars a'_5 and a'_6, a'_7 depending on a_4, a_5 and a_4, a_5, a_6, a_7 resp. By acting $\text{Ad}(\exp(\frac{a'_7}{a'_2} X_4))$ on X' when we suppose that $a_2 \neq 0$, we tend to

$$X'' = a'_1 X_1 + a_2 X_2 + X_3 + a'_5 X_5 + a'_6 X_6$$

where the coefficient of X_5 is vanished for scalar a'_1 depending to a_1, a_2, a'_5, a'_7 . Also be the action of $\text{Ad}(\exp(-(a'_6/a_2) X_5))$ on X'' the coefficient of X_6 will be zero:

$$X''' = a'_1 X_1 + a_2 X_2 + X_3 + a'_5 X_5.$$

At this stage, by acting adjoint representations $M_i(a_i)$ on X''' , we find that no more simplification of X''' is possible. Thus each of a'_1 and a'_5 are arbitrary. Therefore, any one-dimensional Lie subalgebra generated by X with $a_2, a_5 \neq 0$ (by knowing that $a_3 \neq 0$) is equivalent to the Lie subalgebra spanned by

$$a'_1 X_1 + a_2 X_2 + X_3 + a'_5 X_5,$$

which is equal to section 17) of the theorem.

Each of the following cases is prepared by a similar method to *Case 1* and by eliminating unnecessary details, we just give the conditions.

Case 2. If $a_5 \neq 0$ and $a_2 = 0$, then we can make the coefficients of X_6, X_1 vanish resp. by M_2, M_4 and then X is reduced to 11).

Case 3. In the case which $a_5 = 0$ along with $a_2, a_4 \neq 0$, we act on X by M_4, M_5, M_7 to cancel the coefficients of X_7, X_6, X_1, X_2 resp., then the simplest possible form of X is equal to the part 6) of the theorem.

Case 4. When we change the condition $a_4 \neq 0$ in *Case 3.* to $a_4 = 0$, by applying M_4, M_5 the coefficients of X_7, X_6 vanish and no more simplification is possible. Therefore X is reduced to 12).

Case 5. When $a_2 = a_5 = 0$ and $a_4, a_7 \neq 0$, then by applying M_4, M_6, M_2 one can make the coefficients of X_5, X_1, X_7 vanish and hence X turns to 13).

Case 6. If in *Case 5* we only change $a_4 \neq 0$ to $a_4 = 0$ along with $a_6 \neq 0$, then we can act on X by resp. M_4, M_6, M_1 to cancel the coefficients X_5, X_1, X_7 . Hence we find that form of X is in the form of 7).

Case 7. In the latter case, if we suppose that $a_6 = 0$, by acting M_4, M_6 the coefficients of resp. X_5, X_1 are zero and X is equal to the part 8) of the theorem.

Case 8. Let $a_2 = a_5 = a_7 = 0$ and $a_4 \neq 0$. we can make the coefficients of X_5, X_1 resp. by M_4, M_5 vanish and then reduce X to the section 13).

Case 9. When $a_2 = a_4 = a_5 = a_7 = 0$ and $a_6 \neq 0$, by acting resp. M_4, M_7 on X , one can make the coefficients of X_5, X_1 vanish. With these conditions, X gives the form of 7).

Case 10. When $a_2 = a_4 = a_5 = a_6 = a_7 = 0$, by the action of M_4 on X and vanishing X_5 , X is reduced to 5).

• At the second, we suppose that $a_3 = 0$ and consider *Case 11, ..., Case 20* prepared by this assumption.

Case 11. Let $a_4, a_5 \neq 0$. We can act by M_2, M_7, M_4 to cancel the coefficients of X_7, X_2, X_1 resp. then we will have the reduced form 16) of X .

Case 12. By changing $a_5 \neq 0$ of the latter case to be zero and applying M_2, M_7, M_5 on X , the coefficients of X_7, X_2, X_1 resp. will vanish and then X is reduced to 10).

Case 13. Let $a_4 = 0, a_7 \neq 0$ and $a_5 \neq 0$ (by following the process and finding the exact generating coefficients, in fact, we assume that $a'_5 = a_1 a_2 + a_5 a_7$ be non-zero). The coefficients of X_1, X_6 by acting M_6, M_2 will be zero and no further simplification is possible. Hence X is reduced to 15).

Case 14. Suppose that $a_4 = a'_5 = 0, a_7 \neq 0$, and $a_2 \neq 0$ (in fact, in the method of construction, $a'_2 = -a_2 a_7 + a_1 a_5 \neq 0$). Applying M_6, M_4, M_5 resp. on X make the coefficients of X_1, X_7, X_6 vanish. In this case X is reduced to 2).

Case 15. In the last case, we let a'_2 be zero and apply M_6, M_1, M_1 resp. to cancel the coefficients of X_1, X_2, X_6 . Then X will be in the form of 4).

Case 16. If $a'_2 = a_4 = a'_5 = b_7 = 0$ then by acting M_6, M_1 resp. to cancel the coefficients of X_1, X_6 , we lead to the simplest case 3) of X .

Case 17. Let $a_4 = a_7 = 0$ and $a_5 \neq 0$. By acting M_2 on X then the coefficient of X_6 will be zero and hence X is reduced to 14).

Case 18. If $a_4 = a_5 = a_7 = 0$ and $a_2 \neq 0$, then by acting M_5 to cancel the coefficient of X_6 , we lead to the simplest case 9) of X .

Case 19. When $a_2 = a_4 = a_5 = a_7 = 0$ and $a_6 \neq 0$, then we can make the coefficients X_1 resp. by M_7 vanish. Then X is reduced to 3).

Case 18. Finally, if in the last case $a_6 = 0$, no further simplification is possible and then X is reduced to 1).

There is not any more possible case for studying and the proof is complete. $\square$

For finding optimal systems of higher dimensional Lie subalgebras (equivalently Lie subgroups), we refer the reader to [12] or [7].

References

- [1] S. Amini and S.M. Kirkup, *Solution of Helmholtz Equation in the Exterior Domain by Elementary Boundary Integral Methods*, J. Comp. Phys., 118, 208-221 (1995).
- [2] A. Bayliss, C.I. Goldstein, and E. Turkel, *An iterative methods for the Helmholtz equation*, J. Comp. Phys., **49**, 443-457 (1983).
- [3] L.A. Falkovsky and S. Klamazk, *Integral relations and symmetry group expansions for the Helmholtz equation*, J. Phys. A: Math. Gen. **29**, 56275635 (1996).
- [4] N.A. Gumerov and R. Duraiswami, *Fast multipole methods for the Helmholtz equation in three dimensions*, Elsevier, Netherland (2004).
- [5] M.S. Howe, *Acoustics of fluid-structure interactions*, Cambridge Univ. Press (1998).
- [6] N.H. Ibragimov, *Elementary Lie Group Analysis and Ordinary Differential Equations*, John Wiley & Sons, England (1999).
- [7] N.H. Ibragimov, *Selected Works*, Vol. II, Alga Publication, Sweden (2006).
- [8] A. Kushner, V. Lychagin and V. Rubtsove, *Contact Geometry and Non-linear Differential Equations*, Cambridge Univ. Press (2007).
- [9] P. Moon and D.E. Spencer, *Field Theory Handbook, Including Coordinate Systems, Differential Equations, and Their Solutions*, 2nd ed., Springer-Verlag, New York, 15-17 (1988).
- [10] P.J. Olver, *Application of Lie Groups to Differential Equations*, 2nd ed., Springer-Verlag, New York (1993).

- [11] P.J. Olver, *Equivalence, Invariants, and Symmetry*, Cambridge Univ. Press, Cambridge (1995).
- [12] L.V. Ovsiannikov, *Group Analysis of Differential Equations*, Academic Press, New York (1982).